



ANKARA YILDIRIM BEYAZIT
ÜNİVERSİTESİ
TIP FAKÜLTESİ

**ANKARA YILDIRIM BEYAZIT
UNIVERSITY FACULTY OF
MEDICINE**

**2020-2021 EDUCATION YEAR
PHASE I**

**CLINICAL SKILLS TRAINING LEARNING AND
ASSESSMENT GUIDELINES**

BOARD OF CLINICAL SKILLS TRAINING	
HEAD COORDINATOR OF UNDERGRADUATE EDUCATION	PROF. DR. DİDEM ŞENER DEDE
COORDINATOR OF CLINICAL SKILLS EDUCATION	ASSOC. PROF. DR. MEHMET ERGİN
INTERNAL MEDICINE SCIENCES	
DEPT. OF EMERGENCY MEDICINE	ASST. PROF. DR. ÇAĞDAŞ YILDIRIM
DEPT. OF FAMILY MEDICINE	PROF. DR. YUSUF ÜSTÜ
DEPT. OF PEDIATRICS	ASSOC. PROF. DR. SONAY İNCESoy ÖZDEMİR ASSOC. PROF. DR. SARE GÜLFEM ÖZLÜ
DEPT. OF DERMATOLOGY	PROF. DR. AHMET METİN
DEPT. OF INFECTIOUS DISEASES AND CLINICAL MICROBIOLOGY	ASSOC. PROF. DR. BİRCAN KAYAARSLAN
DEPT. OF PHYSICAL MEDICINE AND REHABILITATION	PROF. DR. MÜYESSER ARAS
DEPT. OF CHEST DISEASES	ASSOC. PROF. DR. EMİNE ARGÜDER
DEPT OF INTERNAL MEDICINE	ASSOC. PROF. DR. MUSTAFA TAHTACI (GASTROENTEROLOGY) ASSOC. PROF. DR. BAHAR G. DEMİRCİ (NEPHROLOGY) ASSOC. PROF. DR. ALİ ABBAS TAM (ENDOCRINOLOGY AND METABOLISM DISEASES)
DEPT. OF CARDIOLOGY	ASSOC. PROF. DR. HACI AHMET KASAPKARA
DEPT. OF NEUROLOGY	ASSOC. PROF. DR. MEHMET İLKER YÖN
DEPT. OF PSYCHIATRY	ASSOC. PROF. DR. GÖRKEM KARAKAŞ UĞURLU
SURGICAL MEDICINE SCIENCES	
DEPT. OF ANESTHESIOLOGY AND REANIMATION	ASSOC. PROF. DR. HANDAN GÜLEÇ
DEPT. OF NEUROSURGERY	ASST. PROF. DR. ÜYESİ ERCAN BAL
DEPT. OF GENERAL SURGERY	ASSOC. PROF. DR. ERSİN GÜRKAN DURLU
DEPT. OF OPHTHALMOLOGY	ASSOC. PROF. DR. PINAR NALÇACIOĞLU
DEPT. OF OBSTETRICS AND GYNECOLOGY	ASSOC. PROF. DR. ÜYESİ EMRE ERDEM TAŞ
DEPT. OF CARDIOVASCULAR SURGERY	ASST. PROF. DR. KEMAL EŞREF ERDOĞAN
DEPT. OF OTORHINOLARYNGOLOGY	ASSOC. PROF. DR. KAZIM BOZDEMİR
DEPT. OF ORTHOPEDICS AND TRAUMATOLOGY	ASSOC. PROF. DR. MAHMUT NEDİM AYTEKİN
DEPT. OF PLASTIC RECONSTRUCTIVE AND AESTHETIC SURGERY	PROF. DR. M. EROL DEMİRSEREN
DEPT. OF UROLOGY	ASSOC. PROF. DR. BAHİR GÖK ASST. PROF. DR. ERDEM KOÇ
DEPT. OF CHEST SURGERY	ASST. PROF. DR. İLKNUR ÇELİK

TABLE OF CONTENTS

Rules of Clinical Skills Training	
Evaluation Criteries	
CST Stations	
A. Ability of Applying Basic Life Support for Adults	
B. Ability of Early Intervention During Foreign Body Airway Obstruction	
C. Ability Positioning Patient into Recovery Position	
D. Ability of Hand Washing / Ability of Wearing and Removing the Sterile Gloves	
E. Ability of Measuring Adult Heart Rate (Carodit, Femoral, Radial)	
F. Ability of Measuring Body Temperature (Axillary, Percutaneous, Tympanic)	
G. Ability of Measuring Blood Pressure	
H. Ability of Measuring Respiration Rate / Ability of Measuring Spo2 with a Pulse Oximeter	
I. Ability of Wound Care /First Intervention for Thermal Burn and Medical Dressing	
J. Ability of Immobilizing of Fractures and Dislocations / Ability of Applying Elastic Bandage	
K. Ability of Controlling Bleeding/Epistaxis	
L. Ability of Applying Cervical Collar /Log-rolling and Carrying Injured Person on Trauma Board	

RULES OF CLINICAL SKILLS TRAINING

- 1.** Attendance is compulsory. Related academician will take the roll in each lesson.
- 2.** There will be no change among groups, each student will be ready in the CST hall.
- 3.** Students who come into class after ten mins will be marked absent.
- 4.** Students are supposed to wear white coat and casual outfits that enable them to join practices during CST (for example during life support lesson students will practice chest compression on their knees)
- 5.** During CST lessons mannequins will be treated as if they are real. These lessons are the ones that bring you professional attitudes.
- 6.** Students are expected to be respectful to mannequins and not to hurt them
- 7.** There will be no evaluation at the end of the lesson. Your practical application that you perform after observing instructor will be graded. For this reason, it is important to observe carefully and ask for explanation for the parts you do not understand before it is your turn to practice.

EVALUATION CRITERIA:

It is scored on a scale of 1 to 5.

1. SHOULD BE IMPROVED:

It means unperformed steps or performed incorrectly or not on time.

2. INSUFFICIENT:

It means deficiencies can be found in the performing of many steps.

3. SUFFICIENT:

All the steps can be performed but the deficiencies in the performance mean to require the assistance of the educator.

4. GOOD:

The step is performed appropriately and on time but some steps needed reminders from the educator.

5. EXCELLENT:

The step is performed appropriately and on time without requiring any assistance.

CLINICAL SKILLS TRAINING STATIONS

A, C, J ve L skills will be made in Multidisciplinary Laboratory - II on 1st floor

Other applications will be made in the Clinical Skills Training and Simulation Laboratory on the ground floor.

A. ABILITY OF APPLYING BASIC LIFE SUPPORT FOR ADULTS

SQUAD AND EQUIPMENT:

A half size mannequin, automated external defibrillator training device

PROCESS STEPS:

Basic life support/automated external defibrillator (BLS/AED) algorithm

SEQUENCE/Application	Technical description
SAFETY Make sure you, the victim and any bystanders are safe	
RESPONSE Check the victim for a response	 <p>Gently shake his shoulders and ask loudly: "Are you all right?"</p> <p>If he responds leave him in the position in which you find him, provided there is no further danger; try to find out what is wrong with him and get help if needed; reassess him regularly.</p>
AIRWAY Open the airway	 <p>Turn the patient onto his back if necessary. Place your hand on his forehead and gently tilt his head back; with your fingertips under the point of the victim's chin, lift the chin to open the airway</p>
BREATHING Look, listen and feel for normal breathing	 <p>In the first few minutes after cardiac arrest, a victim may be barely breathing, or taking infrequent, slow and noisy gasps. Do not confuse this with normal breathing. Look, listen and feel for no more than 10 seconds to determine whether the victim is breathing normally. If you have any doubt whether breathing is normal, act as if it is they are not breathing normally and prepare to start CPR.</p>
UNRESPONSIVE AND NOT BREATHING NORMALLY Alert emergency service	 <p>Ask a helper to call the emergency services (112) if possible otherwise call them yourself.</p> <p>Stay with the victim when making the call if possible.</p> <p>Activate speaker function on phone to aid communication with dispatcher</p>
SEND FOR AED Send someone to get AED	 <p>Send someone to find and bring an AED if possible. If you are on your own, do not leave the victim, start CPR.</p>

CIRCULATION

Start chest compressions



Kneel by the side of the victim.

Place the heel of one hand in the centre of the victim's chest; (which is the lower half of the victim's breastbone (sternum))



Place the heel of your other hand on top of the first hand.

Interlock the fingers of your hands and ensure that pressure is not applied over the victim's ribs.

Keep your arms straight.

Do not apply any pressure over the upper abdomen or the bottom end of the bony sternum (breastbone).



Position yourself vertically above the victim's chest and press down the sternum approximately 5 cm (but not more than 6 cm).

After each compression, release all the pressure on the chest without losing contact between your hands and the sternum.

Repeat at the rate of 100-120 per min

IF TRAINED AND ABLE

Combine chest compressions with rescue breaths



After 30 compressions open the airway again using head tilt and chin

Pinch the soft part of the nose closed, using the index finger and thumb of your hand on the forehead.

Allow the mouth to open, but maintain chin lift.

Take a normal breath and place your lips around his mouth, making sure that you have a good seal.

Blow steadily into the mouth while watching for the chest to rise, taking about 1 second as in normal breathing; this is an effective rescue breath.

Maintaining head tilt and chin lift, take your mouth away from the victim and watch for the chest to fall as air comes out.

Take another normal breath and blow into the victim's mouth once more to achieve a total of two effective rescue breaths. Do not interrupt compressions by more than 10 seconds to deliver two breaths. Then return your hands without delay to the correct position on the sternum and give a further 30 chest compressions.

Continue with the chest compressions and rescue breaths in a ratio of 30:2.

IF UNTRAINED OR UNABLE TO DO RESCUE BREATHS

Continue compression only CPR



Give chest compressions only CPR (continuous compressions at the rate of 100-120 per min

WHEN AED ARRIVES

Switch on the AED and attach the electrode pads



As soon as the AED arrives:

Switch on the AED and attach the electrode pads on the victim's bare chest.

If more than one rescuer is present, CPR should be continued while electrode pads are being attached to the chest.

Follow the spoken/visual directions



Ensure that nobody is touching the victim while the AED is analysing the rhythm

If a shock is indicated, deliver shock



Ensure that nobody is touching the victim while AED is analysing the

Push shock buttons as directed (fully automatic AEDs will deliver the shock automatically)

Immediately restart CPR 30:2.

If no shock is indicated, continue CPR



Continue as directed by the voice/visual prompts

Immediately resume CPR. Continue as directed by the voice/visual prompts

**If NO AED IS AVAILABLE
CONTINUE CPR**

Continue CPR



Do not interrupt resuscitation until:

- A health professional tells you to stop
- The victim is definitely waking up moving opening eyes and breathing normally
- You become exhausted

B. ABILITY OF EARLY INTERVENTION DURING FOREIGN BODY AIRWAY OBSTRUCTION

EQUIPMENT: A full size mannequin

PROCEDURE STEPS:

Action	Technical Description
<p>SUSPECT CHOKING</p> <p>Be alert to choking particularly if victim is eating</p> 	
<p>ENCOURAGE TO COUGH</p> <p>Instruct victim to cough</p> 	
<p>GIVE BACK BLOWS</p> <p>If cough becomes ineffective give up to 5 blows</p> 	<p>If the victim shows signs of severe airway obstruction and is conscious apply five back blows. Stand to the side and slightly behind the victim. Support the chest with one hand and lean the victim well forwards so that when the obstructing object is dislodged it comes out of the mouth rather than going further down the airway. Give five sharp blows between the shoulder blades with the heel of your other hand.</p>
<p>GIVE ABDOMINAL THRUSTS</p> <p>If back blows are ineffective give up 5 abdominal thrusts</p> 	<p>If five back blows fail to relieve the airway obstruction, give up to five abdominal thrusts as follows: Stay behind the victim and put both arms round the upper part of the abdomen. Lean the victim forwards. Clench your fist and place it between the umbilicus (navel) and the ribcage. Grasp this hand with your other hand and pull sharply inwards and upwards. If the obstruction is still not relieved, continue alternating five back blows with five abdominal thrusts.</p>
<p>START CPR Start CPR if the victim becomes unresponsive</p> 	<ul style="list-style-type: none"> • If the victim at any time becomes unresponsive: <ul style="list-style-type: none"> <input type="checkbox"/> Support the victim carefully to the ground <input type="checkbox"/> Immediately activate the ambulance service <input type="checkbox"/> Begin CPR with chest compressions.

C. ABILITY OF POSITIONING PATIENT INTO RECOVERY POSITION

EQUIPMENT: Full size mannequin with flexible arms and legs

PROCEDURE STEPS:

1. Lay the person's back, who is unconscious but with breathing and circulation, on a surface.
2. Stand on the right side of the patient.
3. Move the left arm in the position of passing in front of the neck and place the left hand under the right cheek. Position the leg at a right angle (90degrees)
4. Pull the left knee.
5. Pull the patient towards you with your left hand on the shoulder and right hand on hip until the patient's left elbow and left knee touch the floor.
6. Recheck the airway passage, breathing and circulation.
7. If patient does not show any signs of breathing and circulation, apply cardiopulmonary resuscitation by putting his back on a surface.
8. Keep waiting in the same position until ambulance arrives. If it goes on more than 30 min, place the patient' back on the floor and apply the same process for the other person.

IF UNRESPONSIVE BUT BREATHING NORMALLY

If you are certain the victim is breathing normally but is still unresponsive, place in the recovery position



It is rare for CPR alone to restart the heart. Unless you are certain the person has recovered continue

Signs showing the victim recovered:

- Waking up
- moving
- opens eyes
- normal breathing

Be prepared to restart CPR immediately if patient deteriorates.

D. ABILITY OF HAND WASHING

EQUIPMENT: Water, soap/cleaning solution, paper towel, medical waste bag Due to frequent contact with outer media, hands are the most infection causing organs. We ought to know how to wash our hands well and apply it.

PROCEDURE STEPS:

1. Fold the arms of your clothes leaving your wrists open.
2. Remove your jewelry (ring, watch, bracelet).
3. Turn on the tap, if without photocell type.
4. Soak your hands.
5. Get some soap/cleaning solution to your hands.
6. Foam the soap/cleaning solution with some water.
7. If you are using soap, rinse the soap and leave it to its place, when your hands are still foamy.
8. Repeat the following movements for at least five times while your hands are foamy.
 - a. Scrub keeping your palms in touch,
 - b. Scrub the other hand's back with your palm; scrubbing the backside and interdigital areas of the left hand with the palm of the right hand,
 - c. Scrub the back and interdigital areas of the right hand with the palm of the left hand.
 - d. Scrub by passing the fingers through each other.
 - e. For the right and left hands respectively, scrub the backs of the fingers by the other hand's palms.
 - f. Scrub the right and left thumbs respectively, in the other hand's palm.
 - g. Scrub the finger tips of the right and left hand in the other hand's palm.
9. Wash your hands by rubbing and rinse them by shedding the foam of the cleaning solution.
10. Dry your hands with paper towel.
11. While lever faucets are preferred and are turned off with the elbows, classical taps are turned off by pouring water on them from the palms and by holding them with the paper towel used.
12. Throw the used paper towel to the blue household waste bags.

D. ABILITY OF WEARING AND REMOVING STERILE GLOVES

EQUIPMENT: Sterile Gloves

Types of Gloves:

Nylon Gloves: They are made of semi-transparent thin nylon for preventing transmission in food industry. They are loose- fitting. They are not suitable for delicate works.

Examination Gloves: The gloves that are made of latex, are non-sterile and available in standard sizes. They are released in multiple packages, are used frequently, fit the hand.

Sterile Gloves: They are surgical gloves. They are made of latex. The glove shaped for the right hand is different from the one for the left hand. For a better fitting glove, glove sizes range from 7 to 8,5.

PROCEDURE STEPS:

1. Wash and dry your hands according to the instructions.
2. Select proper size of gloves (for sterility).
3. Look for an expiration date on the package of gloves and check gloves for physical damage such as tears.
4. Peel and open the outer package without damaging second paper layer inside.
5. Open the inner package onto the table without touching the gloves and its inside.
6. While putting on, be careful for the place of thumb. Bring the thumb of the glove to the front. Grasp the cuff at the folded edge of the opposing glove with the dominant hand.
7. Slide the other hand into the glove by paying attention to all the fingertips are placed correctly. After fitting your hand into it, pull out the glove to the wrist.
8. While putting on the other hand, pick up the glove by inserting the second and third fingers of the gloved hand into the folded edge. Then, ungloved hand is put on as it is explained above.
9. After completing the process, grasp the cuff of the other glove with your dominant hand and remove it without touching your skin.
10. Throw the removed glove into the medical waste bag (red).
11. Place your ungloved hand between the glove and your wrist without touching the glove exterior. Remove the glove turning it inside out as it goes.

12. Throw the gloves into medical waste bag (red) and the package of the gloves into paper or household waste bags.
13. Wash and dry your hands.

E. ABILITY OF MEASURING ADULT HEART RATE (CAROTID, RADIAL, FEMORAL)

EQUIPMENT: Adult full body mannequin

PROCEDURE STEPS:

1. Wash and dry your hands before and after the examination.
2. Inform the patient about the examination that will be done.
3. Make sure the patient is relaxed and comfortable for 5-10 minutes.
4. Ask him to take off his clothes for watching his chest movements.
5. Stand on the right side of the patient.

MEASURING PULSATION OVER CAROTID

1. Place your index and middle fingers on laryngeal prominence on your neck
2. From this point move your fingers laterally to your side and feel the carotid pulsation on the groove next to the trachea
3. Start chronometer or count the number of beats for one minute looking at the clock
4. Record the number of the beats you have counted in one minute.

MEASURING PULSATION OVER RADIAL ARTERY:

1. Position the wrist, facing the patient's right palm down.
2. Place the tips of the index, middle and ring fingers on the radial artery and palpate.
3. When you feel the pulses, count the number of beats in 1 min
4. Record the number of beats, distension and rhythm.

MEASURING PULSATION OVER FEMORAL ARTERY

1. Lay the patient on his back.
2. Remove his clothes in the manner that his groin area is exposed.
3. The area that you are going to palpate should be just under the inguinal ligament midway between anterosuperior iliac spine and symphysis pubis.
4. Palpate using your 2., 3., 4. fingers.
5. When you feel the pulse count the beats for 1 min.
6. Record the number of beats, distension and rhythm.

F. ABILITY OF MEASURING BODY TEMPERATURE

EQUIPMENT: Axillary, infra-red tympanic thermometer

AXILLARY PROCEDURE STEPS

1. Wash and dry your hands before and after examination.
2. Inform the patient about the examination that will be done.
3. Ask the patient to take off the clothes for watching armpit.
4. Hold the thermometer from its nonmetal part.
5. Turn the thermometer on and check the digital numbers on the screen.
6. Keep the armpit dry.
7. The thermometer is placed under arm with the bulb (metal part) in the center of the armpit.
8. Press the arm against the body and keep the thermometer in the axilla.
9. After waiting for 3 min, take the thermometer out of the axilla and read numbers from the digital screen.
10. Clean the thermometer with a sanitizer and put it away.
11. Provide information to the patient about the measurement results.

PROCEDURE STEPS OF MEASURING TEMPRATURE AT THE SKIN SURFACE

1. Inform the patient about examination that will be done.
2. Wash your hands or use hand sanitizer.
3. Check your equipment.
4. If possible give patient sitting position.
5. Keep forehead and soft depression behind ear dry.
6. Turn on infra-red thermometer.
7. Hold the device 5-10 cm away from the patient.
8. Depress the button and wait until the reading is shown.
9. Record the reading and the area.

TYMPANIC PROCEDURE STEPS

1. Inform the patient about the examination that will be done.
2. Wash your hands or use hand sanitizer
3. Check your equipment
4. If possible, give patient sitting position.
5. Check that there is no plug, discharge or bump that might congest external auditory canal or might prevent measurement.
6. To flatten external auditory canal, pull external ear backwards if the patient is under 1 year old; if the patient is older than 1 year old pull external ear backwards and upwards.
7. Clean the tip of thermometer and put on a new sheath.
8. Turn on thermometer.
9. Place the thermometer probe in external auditory canal.
10. While the thermometer is in external auditory canal depress the button and wait until reading is done.
11. Take thermometer off the ear.
12. Record reading and the area.
13. Throw the used sheath into the bin.

G. ABILITY OF MEASURING BLOOD PRESSURE

EQUIPMENT: Sphygmomanometer

PROCEDURE STEPS:

1. Take patient in a quiet room.
2. Wash your hands before and after the procedure.
3. Inform the patient about the examination that will be done and ask if she/he has smoked, drunk coffee or tea, or done any exercises in 30 mins.
4. Give patient sitting position or help her/him on her/his back (if she/he is sitting feet should be on the ground and uncrossed. Arm should be at heart level.)
5. Slide the clothing on the arm up to shoulder.
6. Let patient relax for 5-10 mins.
7. Place blood pressure cuff 1-2 cm above the antecubital fossa (In order to take blood pressure, the length of the cuff must be $\frac{2}{3}$ length of the upper arm.)
8. Place stethoscope on the brachial artery.
9. Put your hand on the radial artery of the patient and pump the inflation bulb. After radial artery beat is gone, keep inflating for another 30 mmHg.
10. Release open the valve and depressurize the cuff slowly.
11. The first time you hear the Korotkoff sounds clearly is systolic blood pressure and the pressure when Korotkoff sounds stop is diastolic.

H. ABILITY OF MEASURING RESPIRATORY RATE

EQUIPMENT: Breathing mannequin.

PROCEDURE STEPS:

1. Wash your hands before and after the procedure.
2. Inform the patient about the examination that will be done
3. Let patient relax for 5-10 mins.
4. Ask patient to remove her/his clothes in a way you can see chest movements.
5. Stand on the right side of the patient.
6. Place your hand on the patient's chest and observe the chest rise and fall.
7. Count the number of chest movements (while rising) in 1 min and Record.
8. Provide the patient with information about the measurement results.

H. ABILITY OF MEASURING SPO₂ USING PULSE OXIMETER

EQUIPMENT: Pulse oximeter.

PROCEDURE STEPS:

1. Wash your hands before and after the procedure.
2. Inform the patient about the examination that will be done
3. Turn on pulse oximeter
4. Check the finger that you are going to insert into pulse oximeter (There should be no nail polish, henna or paint).
5. Insert the index finger into the pulse oximeter.
6. Wait until pulse oximeter performs measurement.
7. Record the reading.

I. ABILITY OF WOUND CARE & MEDICAL DRESSING

EQUIPMENT: Gauze, water, soap, plaster, bandages.

PROCEDURE STEPS:

1. Wash your hands before and after the procedure.
2. Wear gloves.
3. Inform the patient about the procedure that will be done.
4. Check your supplies.
5. Remove the clothes on wound.
6. Wash the wound are with warm soapy water.
7. Remove small foreign bodies in wound (stones, sand) carefully.
8. Wait for a professional before removing big foreign bodies (knife, glass, etc.) stuck into wound.
9. If there is too much bleeding stop it first. (This is another CSE subject)
10. Clean the sides of the wound with batticon.
11. Evaluate the need for suture.
12. If there is no need for suture and no big foreign body small wound could be left open.
13. For bigger and blistery wounds, it is appropriate to cover the wound with gauze and wrap with bandage (pomade antibiotic is applied between wound and gauze)
14. For small wounds with self-adhesive gauzy tapes can be used.
15. Inform the patient about tetanus vaccine.

I. FIRST INTERVENTION FOR THERMAL BURN AND MEDICAL DRESSING

PROCEDURE STEPS:

1. Wash your hands before and after the procedure
2. Wear gloves
3. Inform the patient about the procedure that will be done
4. Check your supplies
5. Wash the burned area with cold water and stop burning process.
6. Remove the clothes on burned area by cutting them.
7. Rinse the burned surface with warm soapy water.

J. ABILITY OF IMMOBILIZING FRACTURES AND DISLOCATIONS

EQUIPMENT: Cardboard, triangular bandage, adult first aid dummy

PROCEDURE STEPS:

1. When you meet the patient, check the site for safety.
2. Put on your gloves (if there is any).
3. Do not move the deformed and discolored extremity.
4. Place a clean pad on the open wound/bleeding.
5. Splint the deformed and discolored extremity by putting cardboard or wooden board (including upper and lower area) on each sides and stabilize them putting protection pads like cotton or fabric between the skin and the boards.
6. Put 10cm high supporter under the stabilized extremity.
7. In arm injuries, place the arm in the middle of the triangular bandage by protecting its position.
8. Tie the ends of the triangular bandage on back of the neck. After tying as the forearm being elevated at level of the heart, check if the patient has extra problems or not. Then, transfer the patient.

J. ABILITY OF APPLYING ELASTIC BANDAGE

EQUIPMENT: Elastic bandages (in various sizes), adult first aid dummy.

PROCEDURE STEPS:

1. Inform the patient about the examination will be done.
2. Prepare elastic bandages; 16-20cm for lower extremity and 6-10cm for upper extremity.
3. Ask patient to take off the clothes for leaving the extremity to be wrapped completely open.
4. While holding the bandage with your dominant hand, hold the losing end of the bandage with the other hand.
5. By leaving the fingers open on the lower and upper extremities, wrap the bandage around the extremity in a distal to proximal fashion and overlap each turn to cover the previous one by 50 percent.
6. Apply the bandage wrapping cross around the knuckles as the figure of 8.
7. Fasten the losing end to the rest of the bandage.
8. Finally check the circulation on the fingertips (e.g. bruise, pain etc.).

K. ABILITY OF CONTROLLING BLEEDING

EQUIPMENT: Clean gauze pads, bandages, triangular bandages, adult first aid dummy.

PROCEDURE STEPS:

1. When you meet the patient, check the site for safety.
2. Put your gloves on (if there is any).
3. Assist the patient to sit/ lie down.
4. Press on the bleeding area with a clean gauze.
5. Wrap its surface with gauze while pressing.
6. Elevate the bleeding area above the level of heart if it is on the arm or leg.
7. Prepare a triangular bandage for the bleeding on the head.
8. Place a clean gauze on the bleeding area.
9. Place the triangular bandage on the gauze and roll the ends around the head. Then, tie them over the wound.
10. Check if the patient has extra problems or not. Then, transfer the patient.

K. ABILITY OF CONTROLLING EPISTAXIS

PROCEDURE STEPS:

1. When you meet the patient, check the site for safety.
2. Inform the patient about the procedure will be done
3. Put your gloves on (if there is any).
4. Assist the patient to sit/ lie down
5. Give a emesis basin to the patient
6. Squeeze the patients bone and cartilage joint of nose with your thumb and index finger.
7. Continue squeezing for 10 minutes.
8. Release patients nose and check for anterior or posterior bleeding.

L. ABILITY OF APPLYING CERVICAL COLLAR

EQUIPMENT: Adult size cervical collar, adult first aid dummy

PROCEDURE STEPS:

1. When you meet the patient, check the site for safety.
2. Put on your gloves (if there is any).
3. Do not move the patient and his neck.
4. Stabilize the head and the neck of the patient with an assistant.
5. Open the appropriate size collar and place the back of the cervical collar around the patient's neck and make sure that the jaw part is on the front.
6. Place the collar as the front part fitting under the jaw line.
7. Stabilize the collar neither firmly nor loosely and leave the ears outside.
8. Check if the patient has extra problems or not. Then, transfer the patient.

L. ABILITY OF LOGROLLING AND CARRYING INJURED PERSON ON TRAUMA BOARD

EQUIPMENT: Full size first aid mannequin, backboard, 4 practitioners.

PROCEDURE STEPS:

1. When you come across with a traumatized person check the site for safety.
2. Take personal protective precautions.
3. First rescuer, he/she is also the team leader, positions himself/herself at the injured person's head, stabilizes head and shoulders of the patient and coordinates the logroll procedure.
4. Second and third rescuer kneel on the right side of the patient. Second rescuer places his/her hands on the shoulder and hip of the patient, third rescuer places his /her hands around waist and knees of the patient.
5. With the instruction of first rescuer first, second and third rescuers roll the person at the same time.
6. Fourth rescuer positions the backboard underneath the injured person.
7. With the first rescuer's instruction first, second and third rescuer rolls the patient onto the board on his/her back.
8. Afterwards patient is stabilized on the board with belts and foam blocks.
9. Patient is lifted with the instruction of the first rescuer.